scientific reports

OPEN

Reconstructing contact and a potential interbreeding geographical zone between Neanderthals and anatomically modern humans

Saman H. Guran^{1,2,2}, Masoud Yousefi^{3,4}, Anooshe Kafash⁵ & Elham Ghasidian^{2,3}

While the interbreeding of *Homo neanderthalensis* (hereafter Neanderthal) and *Anatomically modern human* (*AMH*) has been proven, owing to the shortage of fossils and absence of appropriate DNA, the timing and geography of their interbreeding are not clearly known. In this study, we applied ecological niche modelling (maximum entropy approach) and GIS to reconstruct the palaeodistribution of Neanderthals and AMHs in Southwest Asia and Southeast Europe and identify their contact and potential interbreeding zone during marine isotope stage 5 (MIS 5), when the second wave of interbreeding occurred. We used climatic variables characterizing the environmental conditions of MIS 5 ca. 120 to 80 kyr (averaged value) along with the topography and coordinates of Neanderthal and modern human archaeological sites to characterize the palaeodistribution of each species. Overlapping the models revealed that the Zagros Mountains were a contact and potential interbreeding zone for the two human species. We believe that the Zagros Mountains acted as a corridor connecting the Palearctic/Afrotropical realms, facilitating northwards dispersal of AMHs and southwards dispersal of Neanderthals during MIS 5. Our analyses are comparable with archaeological and genetic evidence collected during recent decades.

Keywords Zagros Mountains, Ecological niche, Palaeoenvironment, Neanderthals, Anatomically modern humans

Following the ground-breaking discovery of biocultural admixture in the Late Pleistocene of different early human groups of Neanderthals, archaic/modern humans, and Denisovans, a large and growing body of research concerning the nature and evolutionary history of these events is presented. In addition to the signifi ant consequences that the biological exchanges have had on species e.g. and related issues, the time ²⁻⁴, and geography of contact and interbreeding are the subject of intense debate beate. Neanderthals are an extinct lineage of hominins that emerged at approximately 400 kyr and died off at approximately 40 kyr. Fossil localities and morphological evidence of Neanderthals indicate that they are companionable with the Palearctic biogeographical realm, which includes from western Europe to the Altai Mountains in Siberia at 55° latitude and down to approximately 31° in Western Asia ⁷⁻¹⁰. The chronological settlement patterns of the Neanderthals' sites indicate their expansion to the east and southwest Asia from at least 150 kyr¹¹.

On the other hand, Anatomically Modern Humans (AMHs) have evolved in Africa for more than 300 kyr^{12–14}. The evidence, including physical remains and morphological analyses, suggests that they exited Africa over and over during a period of at least 200 kyr^{15–17}. AMHs also reached Eastern Asia at approximately 120 kyr¹⁸ and later reached Europe at approximately 60 kyr^{19,20}. Recent accurate archaeological and palaeoenvironmental data suggest that AMHs rapidly adapted to the new and extreme environments beyond Africa, such as high plateaus, mountain systems and palearctic ecosystems²¹. Moreover, archaeological and fossil evidence indicates that AMHs entered southwestern Asia during MIS 5^{15,22–24}.

¹Institute for Prehistoric Archaeology, University of Cologne, Cologne, Germany. ²DiyarMehr Institute for Palaeolithic Research, Kermanshah, Iran. ³Stiftung Neanderthal Museum, Mettmann, Germany. ⁴Department of Biology, Hakim Sabzevari University, Sabzevar, Iran. ⁵School of Culture and Society, Department of Archaeology and Heritage Studies, Aarhus University, Aarhus, Denmark. [™]email: samanguran@qmail.com

There is strong evidence of multiple interbreeding events between two groups of Neanderthals and archaic/modern humans in western Eurasia e.g. ^{3,25}. Moreover, many attempts have been made to estimate the timing of this interbreeding, and significant success has been achieved e.g. ^{26–28}. Palaeogenetic studies have shown that the second wave of interbreeding occurred during MIS 5^{4,28}. In some studies, researchers have suggested that the lower latitude regions of southwestern Asia have high potential for the first overlap between Neanderthals and AMHs. Sanchez Goñi²⁹ examined the patterns of expansion of Neanderthals and AMHs and showed that they shared the same ecological niches under certain climatic conditions during the Late Pleistocene. Recently, Churchill et al. ³⁰ reported facial morphological similarities between Neanderthals and AMHs in the Near East, indicating it could be a key region for interbreeding between the two lineages. However, it is still unclear where the two species met and interbred.

Ecological niche models (ENMs) are very practical tools for investigating the geography of two species' palaeodistribution and potential interbreeding areas³¹. ENMs have been found to have important applications in palaeobiogeography, archaeology and palaeoanthropology³²-³6. They use occurrence data of a target species, including ancient humans, as well as palaeoenvironmental variables to calculate the probability of a species or hominin species' presence in a defi ed geographic region³7. These models have successfully been used to reconstruct the distribution of different hominin species³5,³6,³8, identify refugia during the ice ages and reconstruct dispersal corridors³9, niche overlap among species³¹ and niche overlap with prey species⁴⁰. For example, Ruan et al.³¹ successfully used ENMs to identify the contact zones of Neanderthals and Denisovans. In another study, Benito et al.³6 applied the ecological niches to determine the distribution of Neanderthals during the last interglacial period in Europe and in the Irano-Turonian region. Thus, ENMs can be used to model the palaeodistribution of Neanderthals and AMHs and locate the geography of their niche overlap³¹,³6.

The aim of the present study was to reconstruct the palaeodistribution of Neanderthals and AMHs during MIS 5 to identify the contact and potential interbreeding geographical zones of these two species. We also estimated the most important predictor of the two species and investigated the responses of the two species to environmental variables. Previous studies have suggested Southwest Asia as a potential area for the interbreeding of Neanderthals and AMHs^{30,41,42}. Notably, this region which is located at the crossroads of the Afrotropical and Palearctic realms⁴³, matches the distribution of AMHs and Neanderthals, respectively. Thus, we hypothesized that these two species fi st met and interbred at the border of these two biogeographic realms where environmental conditions facilitated niche overlap and resource partitioning by providing a highly diverse habitat rich in resources. Climate is a major determinant of species distributions ⁴⁴, particularly at large spatial scales, thus we expect climate to be more effective than topography in shaping the interactions between Neanderthals and AMHs.

Results

Reconstructing the contact and interbreeding zone

The models developed in this study for Neanderthals (AUC = 941) and AMHs (AUC = 895) performed well according to the AUC model performance metric. Our model of the palaeodistribution of Neanderthals shows that north and west of the Mediterranean Sea towards the Levant, vast patches in Turkey, around the Black Sea, south of the Caspian Sea, Taurus, Caucasus and Zagros Mountains, were highly suitable for this species during MIS 5 (Fig. 1). The AMH palaeodistribution model identifi d large and continuous suitable patches in Africa, Arabia and the Iranian Plateau. Our model identifi d the Zagros Mountains as a contact and potential interbreeding zone in Southwest Asia and Southeast Europe.

Variable importance and response curve

We estimated the relative contributions of the environmental variables to the Maxent model of Neanderthals and AMHs. We found that the maximum temperature of the warmest month (with 58.5% contribution), the minimum temperature of the coldest month (with 19.7% contribution), and the annual precipitation (16.5% contribution) were the most important predictors of the palaeodistribution of Neanderthals. The maximum temperature of the warmest month had a negative association with the presence of Neanderthals. Slope (with 35.6% contribution), topographic diversity (with 26% contribution) and precipitation of the warmest quarter (with 14% contribution) were the most important variables in shaping the palaeodistribution of AMHs. Both species presented similar responses to decreases in slope and habitat suitability in areas with high slopes. Figure 2 shows how each environmental variable affects the Maxent prediction for Neanderthals (a) and AMHs (b). The curves show how the predicted probability of presence changes as each environmental variable is varied, keeping all other environmental variables at their average sample value.

Precipitation changes from 140 to 40 kyr

Figure 3 shows the changes in precipitation from 140 to 40 kyr, with 10,000 intervals for the Zagros Mountains. The highest amount of precipitation occurred at 120 kyr, making it a suitable time for range expansion and interactions between Neanderthals and AMHs.

Discussion

Hominin interbreeding is an important topic in palaeoanthropological studies, but when and where it occurred remain largely unknown. Among the different hominin species, the interbreeding of Neanderthals and AMHs is particularly important, as it contributes to the genetics of our own species. Here, we applied ENMs and GIS and revealed that the Zagros Mountains of Iran is a potentially highly suitable geographic unit for niche overlap and a potential interbreeding zone of these two species.




Figure 1. Habitat suitability models of the two *Homo* species and their potential contact and interbreeding zones in Southwest Asia and Southeast Europe. This figure was generated in QGIS 3.14.1 (www.qgis.org). The figures of the Neanderthal (left) and modern human (right) are adapted from www.demorgen.be.

Our niche models predicted niche overlap for the two species in the Zagros Mountains. In support of this fi ding, various studies of genetic data e.g. 45,46, ecological modelling e.g. 21, archaeological and genetic records e.g. 24,46 and fossils 30,47 are in concordance with our niche overlap model. The expansion of Neanderthals to Zagros must have occurred in accordance with the Palearctic environment and karstic terrains from both sides of the Black Sea, i.e., the Caucasus and Anatolia crossing towards the southern regions. The latest evidence has shown that the southernmost expanse of Neanderthals was up to the latitudes of approximately 31° in an arm-shaped area that stretched to the south in two different directions alongside the Anti-Lebanon and Zagros Mountains 9. The Neanderthals in territories further east, such as those found in present-day Uzbekistan 47, Tajikistan 48 and Asian Russia 26,49, are known as Central and North Asian Neanderthals. To date, evidence of the presence of Neanderthals is consistent with southwestern humid mountainous zones, including Anti-Lebanon in the Levant 50, Anatolia 51, the Caucasus 52,53 and Zagros 9,54.

The data on the MP period in the Zagros Mountains region are rich and more up-to-date due to the discoveries of stratifi d sites associated with absolute dates, hominin fossil records, and lithic artefacts. Among the large number of MP sites, four yielded Neanderthal fossils. The best-known of these is Shanidar Cave, where the remains of ten Neanderthals were discovered⁵⁵. Approximately 350 km southeast (approximately 34° latitude), the Wezmeh and Bisetun caves in the Kermanshah region also yielded Neanderthal remains^{9,54,56}. However, the recent discovery of Neanderthal remains from the Bawa Yawan Rock Shelter is signifi ant since it yielded an




Figure 2. Response curves showing how the presence of Neanderthals (a) and AMHs (b) is related to the environmental variables (Bio5: maximum temperature of the warmest month, Bio6: minimum temperature of the coldest month, Bio12: annual precipitation and Bio18: precipitation of the warmest quarter) (https://biodiversityinformatics.amnh.org/open_source/maxent/).




Figure 3. Precipitation changes from 140 to 40 kyr at 10,000 intervals.

in situ Neanderthal tooth in association with the Zagros Mousterian lithic artefacts 9 . The tooth has been dated to around 65 kyr, whereas the age of the Mousterian layer, dates back to 83 kyr 57 .

Owing to the evidence of various hominin fossil remains, it has been determined that the region of Southwest Asia was inhabited by AMH modern humans in the Late Pleistocene. AMHs have inhabited the Levant during at least two periods between 177 and 194 kyr, as evidenced at the site of Misliya¹⁶, and between ~ 120 kyr and

90 kyr, as shown at the sites of Skhul and Qafzeh⁵⁸, before the area was permanently occupied by *H. sapiens* approximately 55,000 years ago⁵⁹. There is a vast amount of data on hominin (including AMHs) occupations from 400 to 50 kyr in Arabia associated with Eastern African lithic technology see ⁴² and references therein, and moreover, physical remains, including AMH finger bones from Al-Wusta dated to ca. 85 kyr¹⁵, all indicate that Arabia was a gateway to Eurasia during the middle to late Pleistocene. There is evidence of the presence of nun-Mousterian MP artefacts dating back to 80 kyr in the southern regions of the Persian Plateau, both in the Zagros⁶⁰ and in the southern to central parts of the Persian Plateau²⁴.

In accordance with our initial expectation, the interaction and potential interbreeding zone of Neanderthals and AMHs was located in the contact zone of the Afrotropical and Palearctic realms, namely, the Zagros Mountains. There are several reasons why the Zagros Mountains are a suitable location for the niche overlap and potential interbreeding zone of two species. First, the Zagros Mountains are characterized by the environmental conditions of the Palearctic realm, which is the birthplace of Neanderthals⁶¹. At the same time, the areas surrounding Zagros are characterized by the environmental conditions of the Afrotropical realm, which is the birthplace of AMHs. Thus, the Zagros Mountains could have been visited repeatedly by people living in the border areas of the Palaearctic and Afrotropical realms during Pleistocene climatic shift . Therefore, the possibility of interactions between different hominins, including AMHs and Neanderthals, was greater in these areas. Second, Zagros covers a vast geographical region (over 1500 km from Lake Van in Turkish Kurdistan to southeastern Iran) capable of supporting large stable human populations. Thi d, Zagros is exceptionally diverse in terms of topography and biodiversity^{40,62-66}, making it capable of supporting the presence of two species at the same time. These mountains facilitate the niche overlap of some animal species with similar niches within the same habitat^{40,67}. These mountains are known to play a very signifi ant role in species distribution by acting as a dispersal barrier or as a dispersal corridor^{63,65}. These findings support the results of our study.

Our fi dings are further supported by new fossil discoveries in the Zagros Mountains and new genetic data⁴⁶. We assume a migration route into the Central Plateau from other directions, including the southern region via Arabia, the Persian Gulf, and the Oman Sea, is plausible⁴⁶. Th s route might have followed the coastal lines towards the north and eventually into the inner parts of the Persian Plateau. Recent evidence of hominin occupations scattered on the surface in areas located in the southernmost part of the Persian Plateau supports our hypothesis^{24,60,68}.

Our initial supposition was that climatic factors would be the predominant force in predicting the distributions of both Neanderthals and AMHs. However, our fi dings revealed a nuanced picture: while climate emerged as the key determinant of the Neanderthal habitat, AMH distribution was signifi antly influenced by topographical variations. The climate was homogenous, but the topography was heterogeneous across the AMH distribution areas. These fi dings likely suggest that topography played a more pronounced role in sculpting the distribution pattern of AMHs. Our study contributes to the growing body of evidence that underscores the complex interplay between environmental factors in determining species distributions. Our results are in line with prey overlap⁴⁰, showing that the annual precipitation and maximum temperature of the warmest month were the most important predictors of Neanderthal distribution on the Persian Plateau. Climate was the most important determinant of Neanderthal distribution in Europe and the Iran–Turanian region during the last interglacial period; however, the influence of topography was confined to local scales³⁶.

One particular application of ecological niche models (ENMs) is to identify suitable areas for the presence of target species where no observations have been made^{69,70}. Field surveys guided by ENMs have led to the discovery of new populations and rare species^{69,70}, thereby proving the utility of ENMs in this context. Our model, which predicts the interbreeding areas of Neanderthals and AMHs, is assigned a very high priority for future fi ld investigations and excavations. Although fi ld testing of ENMs in archaeological studies is limited⁴⁰, we encourage Iranian archaeologists to conduct fi ld excavations in this potential interbreeding area to evaluate the practicality of the models in archaeological research. Moreover, the use of ENMs can guide the allocation of resources for archaeological excavations, ensuring that efforts are concentrated in areas with the highest potential for signifi ant fi dings. By prioritizing these high-probability locations, researchers can maximize the efficicological their fills in the probability locations.


Conclusions

Before this study, our understanding of the interbreeding of AMHs and Neanderthals was based on genetic and morphology data alone 71,72. For the fi st time, we applied ENMs as additional and independent lines of information to locate possible geographic locations where the two species interbred. Our study identifi d the Persian Plateau, particularly the Zagros Mountains, as a potential interbreeding area for AMHs and Neanderthals. The possibility of attracting different hominin groups in the Zagros Mountains is justified by the geographical conditions of this region, since it is located in two different biogeographical zones, namely, the Palearctic and Afrotropical realms. The border areas of two realms are important in biology since they operate as refugia for species from glacial environments. Consequently, some parts of the Zagros Mountains could have been visited repeatedly by people living in the border areas of the Palaearctic and Afrotropical realms during Pleistocene climatic shifts. Therefore, the possibility of interaction between different hominins, including AMHs and Neanderthals, was greater in these areas.

In addition to our fi dings that the Persian Plateau served as a hub for *Homo sapiens* after dispersal from Africa⁴⁶, we conclude that this plateau contributed significantly to hominin distribution^{40,62,73}, dispersal^{24,74,75} and evolution^{46,76}, and we await many exciting discoveries that will shed light on human evolution and dispersal.

Methods Archaeological sites

We obtained 38 occurrence points for Neanderthals and 45 for AMHs (Fig. 4), extracted from multiple sources, including the "Role of Culture in Early Expansions of Humans Out of Africa" (ROCEEH: http://www.roceeh. net) Database (ROAD30,31) and Appendix S1 in Benito et al.³⁶. Each archaeological site was associated with one or two species on the basis of fossil records and lithic artefacts. Since our research focuses on the time frame MIS 5 (e.g., 120–80 kyr), we used only the archaeological sites during this period for southwest Asia and



- 1. Ghar-e Boof
- 2. Jebel Faya FAY-NE1 rock shelter
- 3. Aybut Al Auwa
- 4. Mundafan Al-Buhayrah
- 5.Umm al-Shaa
- 6. Saffagah
- 7. Jebel Qatter 1
- 8. Al Wusta
- 9. Tor Fara
- 10. Nahal Agev
- 11.Abdur

- 12. Gorgora Rock Shelte
- 13. Sodmein Cave
- 14. Kharga Oasis E-76-4
- 15. Nazlet Khater 2
- 16. Derna West
- 17. Hagfet et Tera
- 18. Ksar Akil
- 19. Hayonim Cave
- 20. Tabun Cave
- 21. Hummal Open air
- 22. Dederiyeh cave

- 23. Karain Cave
- 24. Shanidar Cave
- 25. Bawa Yawan Rockshelter
- 26. Qaleh Kurd Cave
- 27. Azokh-1 Cave
- 28.Kudaro III
- 29. Lakonis I
- 30. Krapina
- 31. Grotta Grande di Scario
- 32. Romanelli

Figure 4. Distribution of key archaeological sites dating between MIS 5 to 3 across southwest Asia and southeast Europe. Map data acquired from http://www.roceeh.org and created in www.qgis.org.

southeast Europe. We carefully examined each coordinate and removed the duplicates. Since the environmental data were at a spatial resolution of ~ 5 km (4.65 km at the equator), we thinned the occurrence data to 5 km to avoid pseudoreplication⁴⁰. This time frame was selected because it is suggested that interbreeding events take place during three different periods²⁸. The initial wave of interbreeding occurred ~ 250 to 200 kyr, the second wave of interbreeding occurred ~ 100 to 120 kyr and the third and last interbreeding occurred ~ 60 to 50 kyr. We were unable to find enough archaeological sites associated with the presence of the two species for the fi st interbreeding event to construct robust niche models; thus, we focused on the second interbreeding event that occurred during MIS 5^{28} .

Environmental predictors

We considered environmental variables related to past climate and topography to reconstruct the AMH and Neanderthal niches during MIS 5. As palaeoclimatic variables, we added the maximum temperature of the warmest month, the minimum temperature of the coldest month, the annual precipitation and precipitation of the warmest quarter to the niche models for the MIS 5-time span. Palaeoclimatic data were obtained from Oscillayers, which is a dataset of climatic oscillations over Plio–Pleistocene time scales at high spatial–temporal resolution⁷⁷. We estimated the average values for each of the abovementioned variables during MIS 5 via the raster package v. 3.4–13⁷⁸ implemented in the R environment⁷⁹. To consider topography, we included the slope and topographic heterogeneity^{36,40}, which were downloaded from EarthEnv⁸⁰. To avoid multicollinearity among the predictors, we calculated a variance inflation factor (VIF;⁸¹) via the "vifstep" function in the "usdm" package⁸² and ensured that the collinearity among the predictors was low (VIF < 10).

Ecological niche modelling

In this study, we used the maximum entropy modelling approach⁸³ to reconstruct ecological niche models of Neanderthals and AMHs during MIS 5. Maxent version 3.4.4 was used to build the niche models⁸⁴. We used the Maxent model because it has been shown to perform better than other niche modelling methods^{37,84}. We then overlapped the two palaeodistribution models to identify potential areas for their contact zones in the QGIS (www.qgis.org). The performance of the niche models was assessed via the area under the curve (AUC) metric of the receiving operator characteristic (ROC) curve⁸³. An AUC value of 0.5 indicates that the performance of the model is not better than random, whereas values closer to 1.0 indicate better model performance⁸⁵. The ROC curves were created by selecting 80% of the data for training and 20% for testing.

Data availability

All data needed to evaluate the conclusions in the paper are present in the paper or the references cited here within. We obtained archaeological sites data from the ROCEEH Out of Africa Database (ROAD) (http://www.roceeh.org) and references cited in the manuscript.

Received: 20 April 2024; Accepted: 13 August 2024

Published online: 03 September 2024

References

- 1. Quach, H. et al. Genetic adaptation and neandertal admixture shaped the immune system of human populations. Cell 167(3), 643–656 (2016).
- 2. Green, R. E. et al. A draft sequence of the Neandertal genome. Science 328(5979), 710-722 (2010).
- 3. Prüfer, K. et al. The complete genome sequence of a Nean-derthal from the Altai Mountains. Nature 505(7481), 43-49 (2014).
- 4. Prüfer, K. et al. A high-coverage Neandertal genome from Vindija Cave in Croatia. Science 358(6363), 655-658 (2017).
- 5. Stewart, J. R. & Stringer, C. B. Human evolution out of Africa: The role of Refugia and climate change. *Science* 335(6074), 1317–1321 (2012).
- 6. Higham, T. et al. The timing and spatiotemporal patterning of Neanderthal disappearance. Nature 512(7514), 306–309 (2014).
- 7. Groucutt, H. S. et al. Rethinking the dispersal of Homo sapiens out of Africa. Evolution Anthropol Issues News Rev 24(4), 149–164 (2015).
- Kolobova, K. A. et al. Archaeological evidence for two separate dispersals of Neanderthals into southern Siberia. Proceedings of the National Academy of Sciences 117(6), 2879–2885 (2020).
- 9. Heydari-Guran, S. et al. The discovery of an in situ Neanderthal remain in the Bawa Yawan rockshelter, west-central Zagros Mountains. Kermanshah. PloS one 16(8), e0253708 (2021).
- 10. Been, E. et al. The first Neanderthal remains from an open-air Middle Palaeolithic site in the Levant. Scientific Reports 7(1), 2958 (2017).
- 11. Grün, R. & Stringer, C. Tabun revisited: revised ESR chronology and new ESR and U-series analyses of dental material from Tabun C1. *J Human Evol* **39**(6), 601–612 (2000).
- 12. Hublin, J. J. et al. New fossils from Jebel Irhoud, Morocco and the pan-African origin of Homo sapiens. Nature **546**(7657), 289–292 (2017).
- 13. Richter, D. et al. The age of the hominin fossils from Jebel Irhoud, Morocco, and the origins of the Middle Stone Age. Nature 546(7657), 293–296 (2017).
- 14. Stringer, C. & Galway-Witham, J. On the origin of our species. Nature 546(7657), 212-214 (2017).
- 15. Groucutt, H. S. et al. Homo sapiens in Arabia by 85,000 years ago. Nat Ecol Evol 2(5), 800-809 (2018).
- 16. Hershkovitz, I. et al. The earliest modern humans outside Africa. Science 359(6374), 456-459 (2018).
- 17. Harvati, K. et al. Apidima Cave fossils provide earliest evidence of Homo sapiens in Eurasia. Nature 571(7766), 500-504 (2019).
- 18. Liu, W. et al. Human remains from Zhirendong, South China, and modern human emergence in East Asia. Proceedings of the National Academy of Sciences 107(45), 19201–19206 (2010).
- 19. Slimak, L. et al. Modern human incursion into Nean-derthal territories 54,000 years ago at Mandrin France. Sci. Adv. 8(6), 9496
- 20. Hublin, J. J. et al. Initial upper palaeolithic homo sapiens from bacho kiro cave Bulgaria. Nature 581(7808), 299-302 (2020).
- 21. Roberts, P. & Stewart, B. A. Defini g the 'generalist specialist'niche for Pleistocene Homo sapiens. *Nat. Human Behav.* **2**(8), 542–550 (2018).

- 22. Groucutt, H. S. & Petraglia, M. D. The prehistory of the Arabian peninsula: Deserts, dispersals, and demography. Evol. Anthropol. Issues News Rev. 21(3), 113–125 (2012).
- Scerri, E. M. et al. Middle to Late Pleistocene human habitation in the western Nefud desert, Saudi Arabia. Quarter. Int. 382, 200–214 (2015).
- 24. Heydari-Guran, S. & Ghasidian, E. (2021). Consistency of the 'MIS 5 Humid Corridor Model' for the Dispersal of Early Homo sapiens into the Iranian Plateau. In: Pearls, Politics and Pistachios: Essays in Anthropology and Memories on the Occasion of Susan Pollock's 65th Birthday 219–238.
- 25. Slon, V. et al. The genome of the offspring of a Neanderthal mother and a Denisovan father. Nature 561(7721), 113-116 (2018).
- Krause, J. et al. The complete mitochondrial DNA genome of an unknown hominin from southern Siberia. Nature 464(7290), 894–897 (2010).
- 27. Yuan, K. et al. Refining models of archaic admixture in Eurasia with ArchaicSeeker 2.0. Nat. Commun. 12(1), 6232 (2021).
- 28. Li, L., Comi, T. J., Bierman, R. F. & Akey, J. M. Recurrent gene fl w between Neanderthals and modern humans over the past 200,000 years. *Science* 385(6705), eadi1768 (2024).
- 29. Goñi, M. F. S. Regional impacts of climate change and its relevance to human evolution. Evolutionary Human Sciences 2, e55 (2020).
- 30. Churchill, S. E., Keys, K. & Ross, A. H. Midfacial Morphology and Neandertal-Modern Human Interbreeding. *Biology* 11(8), 1163 (2022)
- 31. Ruan, J. et al. Climate shifts orchestrated hominin interbreeding events across Eurasia. Science 381 (6658), 699-704 (2023).
- Varela, S., Lobo, J. M. & Hortal, J. Using species distribution models in paleobiogeography: a matter of data, predictors and concepts. Palaeogeogr. Palaeoclimatol. Palaeoecol. 310, 451–463 (2011).
- 33. Svenning, J. C., Fløjgaard, C., Marske, K. A., Nógues-Bravo, D. & Normand, S. Applications of species distribution modeling to paleobiology. *Quat. Sci. Rev.* 30, 2930–2947 (2011).
- Franklin, J., Potts, A. J., Fisher, E. C., Cowling, R. M. & Marean, C. M. Paleodistribution modeling in archaeology and paleoanthropology. *Quat. Sci. Rev.* 110, 1–14 (2015).
- 35. Giampoudakis, K. *et al.* Niche dynamics of Palaeolithic modern humans during the settlement of the Palaearctic. *Global Ecol. Biogeogr.* **26**(3), 359–370 (2017).
- 36. Benito, B. M. et al. The ecological niche and distribution of neanderthals during the last interglacial. J. Biogeogr. 44, 51-61 (2017).
- 37. Guisan, A., Thuiller, W. & Zimmermann, N. E. Habitat Suitability and Distribution Models: With Applications in R (Cambridge University Press, 2017).
- Jones, P. J., Williamson, G. J., Bowman, D. M. & Lefroy, E. C. Mapping Tasmania's cultural landscapes: Using habitat suitability modelling of archaeological sites as a landscape history tool. J. Biogeogr. 46(11), 2570–2582 (2019).
- 39. Kafash, A., Yousefi, M., Ghasidian, E., & Heydari-Guran, Saman (2023). Reconstruction of Epipaleolithic settlement and "Climatic Refugia" in the Zagros Mountains During the Last Glacial Maximum (Lgm). Archeologia e Calcolatori, 34(1).
- 40. Yousefi, M., Heydari-Guran, S., Kafash, A. & Ghasidian, E. Species distribution models advance our knowledge of the Neanderthals' paleoecology on the Iranian Plateau. *Sci Rep* 10, 14248 (2020).
- 41. Groucutt, H. S., Scerri, E. M., Stringer, C. & Petraglia, M. D. Skhul lithic technology and the dispersal of Homo sapiens into Southwest Asia. *Quaternary International* 515, 30–52 (2019).
- 42. Groucutt, H. S. et al. Multiple hominin dispersals into Southwest Asia over the past 400,000 years. Nature 597(7876), 376–380 (2021).
- 43. Holt, B. G. et al. An update of Wallace's zoogeographic regions of the world. Science 339(6115), 74-78 (2013).
- 44. Sexton, J. P., McIntyre, P. J., Angert, A. L. & Rice, K. J. Evolution and ecology of species range limits. *Annu. Rev. Ecol. Evol. Syst.* 40, 415–436 (2009).
- 45. Lazaridis, I., Belfer-Cohen, A., Mallick, S., Patterson, N., Cheronet, O., Rohland, N., & Reich, D. (2018). Paleolithic DNA from the Caucasus reveals core of West Eurasian ancestry. BioRxiv, 423079.
- Vallini, L. et al. The Persian plateau served as hub for Homo sapiens after the main out of Africa dispersal. Nat. Commun. 15(1), 1882 (2024).
- 47. Glantz, M. et al. New hominin remains from Uzbekistan. J. Human Evol. 55(2), 223-237 (2008).
- 48. Trinkaus, E., Ranov, V. A. & Lauklin, S. Middle Paleolithic human deciduous incisor from Khudji Tajikistan. *J. Human Evol.* 38(4), 575–584 (2000).
- 49. Mednikova, M. B. (2014). Neanderthal presence in southern Siberia on data of postcranial morphology. Cultural developments in Eurasian Paleolithic and the origin of anatomically modern humans. –Novosibirsk: Inst. of archaeol. and ethnogr. SB RAS, 158–164.
- 50. Tillier, A. M., & Arensburg, B. (2016). Neanderthals and modern humans in the Levant: An Overview. Quaternary environments, climate change and human in the Levant, 607–610.
- 51. Harvati, K., & Roksandic, M. (Eds.). (2017). Paleoanthropology of the Balkans and Anatolia: Human evolution and its context. Springer.
- 52. Hajdinjak, M. et al. Reconstructing the genetic history of late Neanderthals. Nature 555(7698), 652–656 (2018).
- 53. Gasparyan, B., & Glauberman, P. (2022). Beyond European boundaries: Neanderthals in the Armenian Highlands and the Caucasus. In Updating Neanderthals (pp. 275–301). Academic Press.
- Zanolli, C. et al. A Neanderthal from the Central Western Zagros, Iran. Structural reassessment of the Wezmeh 1 maxillary premolar. J. Human Evol. 135, 102643 (2019).
- 55. Pomeroy, E. et al. New Neanderthal remains associated with the 'fl wer burial' at Shanidar Cave. Antiquity 94(373), 11-26 (2020).
- 56. Trinkaus, E., & Biglari, F. (2006). Middle paleolithic human remains from Bisitun cave, Iran. Paléorient, 105-111.
- 57. Heydari, M., Guérin, G. & Heydari-Guran, S. A Bayesian luminescence chronology for the Bawa Yawan Rock Shelter at the Central Zagros Mountains (Western Iran). *Quaternary International* **680**, 64–75 (2024).
- 58. Mercier, N. et al. Thermoluminescence date for the Mousterian burial site of Es-Skhul Mt. Carmel. J. Archaeol. Sci. 20(2), 169–174 (1993).
- 59. Hershkovitz, I. *et al.* Levantine cranium from Manot Cave (Israel) foreshadows the fi st European modern humans. *Nature* **520**(7546), 216–219 (2015).
- 60. Heydari, M., Guérin, G., Zeidi, M. & Conard, N. J. Bayesian luminescence dating at Ghār-e Boof, Iran, provides a new chronology for Middle and Upper Paleolithic in the southern Zagros. *J. Human Evol.* **151**, 102926 (2021).
- 61. Harvati, K. (2007). Neanderthals and their contemporaries. In Handbook of paleoanthropology (pp. 1717–1748). Springer.
- 62. Heydari-Guran, S. Palaeolithic landscapes of Iran S2586 (BAR Publishing, 2014).
- 63. Kafash, A. et al. Reptile species richness associated to ecological and historical variables in Iran. Sci. Rep. 10, 18167 (2020).
- 64. Kafash, A., Ashrafi, S. & Yousefi, M. Biogeography of bats in Iran: Mapping and disentangling environmental and historical drivers of bat richness. *J. Zool. Syst. Evol. Res.* **59**, 1546–1556 (2021).
- Yousefi, M., Mahmoudi, A., Vaissi, S. & Kafash, A. Diversity, diversification and distribution of Iranian vertebrates: the legacy of mountains uplifting, past climatic oscillations, sea level fluctuations and geographical barriers. *Biodivers Conserv* 32, 7–36 (2023).
- 66. Yousefi, M., Mahmoudi, A., Kafash, A., Khani, A. & Kryštufek, B. Biogeography of rodents in Iran: species richness, elevational distribution and their environmental correlates. *Mammalia* 86, 309–320 (2022).
- 67. Grant, P. The classical case of character displacement. Evol. Biol. 8, 237-337 (1975).

- 68. Shoaee, M. J., Nasab, H. V. & Petraglia, M. D. The Paleolithic of the Iranian Plateau: Hominin occupation history and implications for human dispersals across southern Asia. *J. Anthropol. Archaeol.* **62**, 101292 (2021).
- 69. Fois, M., Fenu, G., Cuena Lombraña, A., Cogoni, D. & Bacchetta, G. A Practical method to speed up the discovery of unknown populations using species distribution models. *J. Nat. Conserv.* 24, 42–48 (2015).
- 70. Becker, F. S., Slingsby, J. A., Measey, J., Tolley, K. A. & Altwegg, R. Finding rare species and estimating the probability that all occupied sites have been found. *Ecol. Appl.* 32, e2502 (2022).
- 71. Trinkaus, E. Modern human versus neandertal evolutionary distinctiveness. Curr. Anthropol. 47(4), 597-620 (2006).
- 72. Tobler, R. et al. The role of genetic selection and climatic factors in the dispersal of anatomically modern humans out of Africa. Proceedings of the National Academy of Sciences 120(22), e2213061120 (2023).
- 73. Vahdati Nasab, H. et al. Qaleh Kurd Cave (Qazvin, Iran): Oldest evidence of middle pleistocene hominin occupations and a human deciduous tooth in the Iranian central plateau. J. Paleolithic Archaeol. 7(1), 1–28 (2024).
- 74. Ghasidian, E., Kafash, A., Kehl, M., Yousefi, M. & Heydari-Guran, S. Modelling Neanderthals' dispersal routes from Caucasus towards east. *PLoS One* 18, e0281978 (2023).
- 75. Shoaee, M. J. et al. Defini g paleoclimatic routes and opportunities for hominin dispersals across Iran. PLoS One 18, e0281872 (2023)
- 76. Ghasidian, E., Heydari-Guran, S. & Mirazón Lahr, M. Upper Paleolithic cultural diversity in the Iranian Zagros Mountains and the expansion of modern humans into Eurasia. *J. Hum. Evol.* **132**, 101–118 (2019).
- 77. Gamisch, A. Oscillayers: A dataset for the study of climatic oscillations over Plio-Pleistocene time-scales at high spatial-temporal resolution. *Global Ecol. Biogeogr.* **28**, 1552–1560 (2019).
- 78. Hijmans, R. J. (2021). raster: Geographic Data Analysis and Modeling. R package version 3.4-13.
- 79. R Core Team. R: A Language and Environment for Statistical Computing (R Foundation for Statistical Computing, 2023).
- 80. Amatulli, G. et al. A suite of global, cross-scale topographic variables for environmental and biodiversity modeling. Scientific Data 5, 180040 (2018).
- 81. Quinn, G. P. & Keough, M. J. Experimental design and data analysis for biologists (Cambridge University Press, 2002).
- 82. Naimi, B. (2015). Uncertainty analysis for species distribution models. R package version 1.1-15.
- 83. Phillips, S. J., Anderson, R. P. & Schapire, R. E. Maximum entropy modeling of species geographic distributions. *Ecological Modeling* 190, 231–259 (2006).
- 84. Elith, J. et al. Novel methods improve prediction of species' distributions from occurrence data. Ecography 29, 129-151 (2006).
- 85. Swets, J. A. Measuring the accuracy of diagnostic systems. Science 240, 1285-1293 (1988).

Acknowledgements

All data needed to evaluate the conclusions in the paper are present in the paper or the references cited here within. We obtained archaeological sites data from the ROCEEH Out of Africa Database (ROAD) (http://www.roceeh.org) and references cited in the manuscript.

Author contributions

"S.H.G., M.Y., and contributed equally to the design and conceptualization and writing and A.K. in performance and analysis of the data. E. G. in review & editing. All authors reviewed the manuscript."

Funding

Open Access funding enabled and organized by Projekt DEAL. The research was funded by the Deutsche Forschungsgemeinschaft (DFG), Priority program 2176: The Iranian highlands: Resiliencies and integration in premodern societies, "Last Neanderthal and early *Homo sapiens* occupations in the Bawa Yawan Rockshelter, Kermanshah, West-Central Zagros Mountains (Iran), under project number 402379177.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to S.H.G.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

© The Author(s) 2024